

## Biosorption of cobalt (II) by Intact and Chemically Modified Brown Algae: Optimization Using Response Surface Methodology and Equilibrium, Dynamics and Thermodynamics Studies

F. Soleymani<sup>1</sup>, H. Pahlevanzadeh<sup>1\*</sup>, M.H. Khani<sup>2</sup>, M. Manteghian<sup>1</sup>

1- Chemical Engineering Department, Tarbiat Modares University, Tehran, Iran

2- Nuclear Fuel Cycle Research School, Nuclear Science and Technology Research Institute, Tehran, Iran

### Abstract

By using response surface methodology, Batch shaking biosorption of cobalt (II) experiments were conducted in order to examine the combined effects of operating parameters. The results indicate that magnesium nitrate performed as an effective biosorbent surface modifier, which increases the rate of adsorption capacity. At optimal conditions (initial pH 7.0, temperature 45°C, biosorbent concentration 0.1 g/100ml, and initial cobalt concentration 300mg/l for Mg-treated biomass) the biosorption capacity of the algae for cobalt was found to be 80.55 mg/g. The Langmuir and Freundlich isotherms were applied to the equilibrium data. The results are best fitted by the Freundlich model. Evaluation of the experimental data in terms of biosorption dynamics showed that the biosorption of cobalt (II) onto algal biomass followed the pseudo-second-order dynamics well. Using the thermodynamic equilibrium coefficients obtained at different temperatures, the thermodynamic parameters ( $\Delta G^\circ$ ,  $\Delta H^\circ$  and  $\Delta S^\circ$ ) were also evaluated.

**Keywords:** Biosorption, Response Surface Methodology, Pretreated Algae, Cobalt, *Sargassum* sp.

### 1. Introduction

Cobalt containing compounds discharged through different industrial activities constitute one of the major causes of water pollution. Heavy metals pose a significant threat to the environment and public health because of their toxicity, accumulation in the food chain and persistence in nature [1,2]. So it is necessary to remove cobalt at the time of release of effluent. There are some conventional methods for treatment of liquid effluent for removal of heavy metals such as

ion exchange, chemical precipitation, reverse osmosis, and co-precipitation which are designed to remove radionuclides and metal ions from the effluents [3]. These conventional techniques can reduce metal ions, but they do not appear to be highly effective due to the limitations in the pH range as well as the high material and operational costs [4]. Therefore, there is growing interest in using low-cost, easily available materials for the adsorption of metal ions. A low-cost adsorbent is defined as one which is abundant in nature, or is a by-product or waste material from another

---

\*Corresponding author: pahlavzh@modares.ac.ir

industry. Biosorbent materials derived from suitable biomass can be used for the effective removal and recovery of heavy metal ions from industrial solutions. Many algae, yeasts, bacteria and other fungi are known to be capable of concentrating metal species from dilute aqueous solutions and accumulating them within their cell structure [5]. Among biosorbent materials algae have proved to be both economic and eco-friendly as they are abundantly available, have regeneration and metal recovery potentiality, lesser volume of chemical and/or biological sludge, high efficiency in dilute effluents, and a high surface area to volume ratio [6]. The cell walls of brown algae generally contain components such as cellulose and alginic acid which contain several functional groups (amino, carboxyl, sulphate and hydroxyl) that could play an important role in the biosorption process [7]. The binding strength of alkaline earth metals to alginic acid was found to decrease in the order Ba(II) > Sr (II) > Ca (II) > Mg (II) [8,9]. Haug interpreted the preferential binding of heavier ions to stereochemical effects, since larger ions might better fit a binding site with two distant functional groups [10]. Also, the preferential binding of larger ions can be attributed to stereochemical effects such as the coordination of the oxygen atoms surrounding the metal ion [11]. According to the trends, cobalt ion is preferable to replace magnesium instead of calcium ion because Mg(II) has less binding strength. So for pretreatment of biomass, Mg(II) was used as a surface modifier instead of other ions such as Ca(II).

The aim of this study was to conduct a central composite design (CCD) analysis to

find the significant factors that influenced the removal of cobalt from aqueous solution by marine brown algae *Sargassum sp.* and understand their impact on the process. The effect of some operating variables such as chemical modification of the biomass, temperature, pH, adsorbent dose and initial concentration of  $\text{Co}^{2+}$  on biosorption was studied using CCD, which gives a mathematical model that shows the influence of each variable and their interactions. In addition, the equilibrium, dynamic and thermodynamic parameters were calculated. In the statistical design, the factors involved in an experiment are simultaneously changed. The most important advantages are that not only the effects of individual parameters but also their relative importance in a given process are evaluated and the interaction of two or more variables can also be derived. This is not possible in a classical one factor at a time experiment [2].

## 2. Mathematical description

The amount of metal uptake by algae was calculated from the differences between the metal quantity added to the biomass and the metal content of the supernatant using the following equation:

$$q = \frac{V(C_0 - C)}{M_{\text{ads}}} \quad (1)$$

where  $q$  is the amount of metal ions adsorbed on the biosorbent at any time,  $t$ , in mg/g,  $V$  is the volume of metal containing solution in contact with the biosorbent in ml,  $C_0$  is the initial concentration of metal in mg/l,  $C$  is the metal concentration in the solution at any

time  $t$  (min) in mg/l, and  $M_{ads}$  is the amount of added biosorbent on dry basis in g [12].

### 2-1. Dynamic parameters of biosorption

It is known that the biosorption process is time-dependent; so in this study the pseudo-first-order and pseudo-second-order dynamic models have been discussed to identify the rate and kinetics of sorption of cobalt on *Sargassum sp.* adsorbent.

Pseudo-first-order Lagergren model considers that the rate of occupation of biosorption sites is proportional to the number of unoccupied sites. The linear form of the model is:

$$\log(q_{eq} - q_t) = \log q_{eq} - \frac{k_1 t}{2.303} \quad (2)$$

where  $q_{eq}$  and  $q_t$  ( $\text{mg g}^{-1}$ ) are amount of cobalt adsorbed at equilibrium and at time  $t$  (min), respectively, and  $k_1$  ( $\text{min}^{-1}$ ) shows the equilibrium rate constant of pseudo-first-order adsorption.  $k_1$  and  $q_{eq}$  can be determined from the linear plot of  $\log(q_{eq} - q_t)$  versus  $t$ .

The adsorption dynamics may also be described by pseudo-second-order model. Pseudo-second-order model considers that the rate of occupation of biosorption sites is proportional to the square of the number of unoccupied sites and after being integrated and rearranged:

$$\frac{t}{q_t} = \frac{1}{k_2 q_{eq}^2} + \frac{1}{q_{eq}} t \quad (3)$$

Where  $q_{eq}$  and  $q_t$  ( $\text{mg g}^{-1}$ ) are amount of cobalt adsorbed at equilibrium and at time  $t$  (min), respectively, and  $k_2$  ( $\text{g mg}^{-1} \text{min}^{-1}$ ) is

the equilibrium rate constant of pseudo-first-order adsorption.  $k_2$  and  $q_{eq}$  can be determined from the linear plot of  $t/q_t$  versus  $t$ . This model is more likely to predict dynamic behavior of biosorption with chemical sorption being the rate-controlling step. Due to the presence of greater number of metal ions in industrial wastewater, the sorption equilibrium was reached much faster due to faster occupancy of sorption sites by metal ions [2,13].

### 2-2. Equilibrium parameters of biosorption

The biosorption isotherms are characterized by definite parameters whose values express the surface properties and affinity of biosorbent for different heavy metal ions [14]. Several isotherm equations have been used for the equilibrium modeling of biosorption systems. In this study, the Langmuir and Freundlich adsorption models were tested to determine the relationship between adsorbed cobalt ions on the algal cell ( $q_{eq}$ ) and unadsorbed cobalt ions in solution ( $C_{eq}$ ). The most widely used isotherm equation for modeling equilibrium is the Langmuir equation, based on the assumption that there is a finite number of binding sites which are homogeneously distributed over the adsorbent surface, these binding sites have the same affinity for adsorption of a single molecular layer and there is no interaction between adsorbed molecules [5]. The linear form of Langmuir isotherm is given by the following equation:

$$\frac{1}{q_{eq}} = \frac{1}{b q_m C_{eq}} + \frac{1}{q_m} \quad (4)$$

where  $q_{eq}$  ( $\text{mg/g}$ ) is the amount adsorbed at

the equilibrium,  $C_{eq}$  (mg/l) the equilibrium concentration,  $q_m$  (mg/g) is the Langmuir constant representing the maximum monolayer adsorption capacity and  $b$  (l/mg) is the Langmuir constant related to energy of adsorption and the affinity of the binding sites [1,15]. The equilibrium parameters ( $q_m$  and  $b$ ) are determined from the linear plot of  $1/q_{eq}$  versus  $1/C_{eq}$ .

The adsorption data was also analyzed by Freundlich model. The Freundlich expression is used for heterogeneous surface energy term. The Freundlich isotherm equation is an exponential equation and therefore, assumes that as the adsorbate concentration increases, the concentration of adsorbate on the adsorbent surface also increases [5]. Freundlich equation is commonly presented as shown below:

$$q_{eq} = kC_{eq}^{\frac{1}{n}} \quad (5)$$

The logarithmic form of Freundlich model is given by the following equation:

$$\log q_{eq} = \log k + \frac{1}{n} \log C_{eq} \quad (6)$$

where  $k$  and  $n$  are constants related to the adsorption capacity and intensity of the adsorbent characteristics of the system, respectively. The fractional values of biosorbent surface indicate a favorable biosorption of metal ions onto biomass [16]. The plots of  $\log q_{eq}$  versus  $C_{eq}$  determine values of  $1/n$  and  $k$ .

### 2-3. Thermodynamic parameters of biosorption

The determination of thermodynamics parameters has great importance in

evaluating spontaneity and heat change for the biosorption reactions. The Gibbs free energy change of the adsorption process ( $\Delta G^\circ$ ) is related to the standard thermodynamic equilibrium constant ( $K_{eq}^\circ$ ) of the biosorption system by the classic equation [17,18]:

$$\Delta G^\circ = -RT \ln K_{eq}^\circ \quad (7)$$

Where  $T$  is the absolute temperature (kelvin) and  $R$  is the universal gas constant which is 8.314 (j/mol k).

The apparent equilibrium constant ( $K'_e$ ) of the adsorption is defined as [17, 19]:

$$K'_e = \frac{C_{ad,eq}}{C_{eq}} \quad (8)$$

Where  $C_{ad,eq}$  is the amount of adsorbed metal ion on the biosorbent per volume of the solution at equilibrium and  $C_{eq}$  is the equilibrium concentration of the metal ion in the solution. The Gibbs free energy change ( $\Delta G^\circ$ ) reflects the feasibility and spontaneous nature of the process. At a given temperature, the higher the negative value of  $\Delta G^\circ$ , the more energetically favorable adsorption process.

In order to determine the nature of the biosorption process regarding heat of adsorption and also affinity of biosorbent to metal ion, enthalpy changes ( $\Delta H^\circ$ ), and entropy changes ( $\Delta S^\circ$ ) should also be estimated. These parameters can be estimated by the van't Hoff equation [17, 19]:

$$\ln K_{eq}^\circ = \frac{\Delta S^\circ}{R} - \frac{\Delta H^\circ}{RT} \quad (9)$$

Where  $\Delta H^\circ$  and  $\Delta S^\circ$  can be obtained from the slope and intercept of a van't Hoff plot of  $\ln K^\circ$  vs  $1/T$ .

### **3. Materials and methods**

#### **3-1. Preparation of biosorbents**

Samples of marine algae *Sargassum sp.* were collected from Persian Gulf on the coast of Qeshm, Iran. Algal samples were washed with tap water and distilled water to remove sand and excess sodium and potassium ions. After drying overnight at a maximum temperature of 55°C to avoid degradation of the binding sites [14], the samples were grounded to an average particle size of 0.7 mm. The biomasses were subsequently loaded with Mg<sup>2+</sup> in a solution of 0.1 M Mg(NO<sub>3</sub>)<sub>2</sub>.6H<sub>2</sub>O (biomass concentration of 10 g/l) for 24 h under slow stirring. Then pretreated biomass was washed several times with deionized water until a stable wash solution pH was reached and excess magnesium ions were removed from the biomasses. The Mg-pretreated biomasses were then dried overnight in an oven at 55°C and subsequently natural algae and Mg-pretreated algae were used for biosorption experiments.

#### **3-2. Preparation of synthetic metal solution**

A stock solution of cobalt, with 1000 ppm concentration was prepared using analytical grade (Co(NO<sub>3</sub>)<sub>2</sub>.6H<sub>2</sub>O). This solution was diluted depending upon requirements. All solutions were prepared in deionized water. The initial pH of cobalt solutions was adjusted with diluted or concentrated HNO<sub>3</sub> and NaOH solutions before mixing with the biosorbent. All chemicals used were of analytical grade (Sigma Aldrich, Germany).

#### **3-3. Characterization test**

##### **3-3-1. Energy dispersive X-ray spectroscopy (EDX)**

To determine the chemical composition of biosorbent before and after treatment by 0.1

M Mg(NO<sub>3</sub>)<sub>2</sub>.6H<sub>2</sub>O and also after Co<sup>2+</sup> sorption, a system of analysis for energy dispersive X-ray spectroscopy (EDX, RONTECH, Germany) was used.

##### **3-3-2. Analysis of cobalt concentration**

Dissolved cobalt concentrations in solution were assessed by an inductively coupled plasma atomic emission spectrophotometer (ICP-AES, Optima, 7300DV, USA). The ICP analyses were conducted at wavelength of 228.616 nm.

##### **3-4. Batch adsorption experiments**

In order to determine the contact time required for the sorption equilibrium experiments, the sorption dynamics experiments were conducted on natural and Mg-pretreated algae first. 0.1 gram of biomass was added to a series of 250 ml Erlenmeyer flasks containing 100 ml of 100 mg Co/l Co(NO<sub>3</sub>)<sub>2</sub>.6H<sub>2</sub>O solution. The flasks were agitated at 150 rpm and 25°C for 24 h in a shaker. Samples were withdrawn at predetermined time intervals (2, 5, 15, 30, 45, 60, 90, 120, 150, 180 and 1440 min). After appropriate dilution, the samples were analyzed by the ICP-AES for metal concentrations. According to the preliminary sorption dynamic tests, the equilibrium was reached after 90 min of contact. Batch adsorption experiments were conducted to study the effect of magnesium ion as a surface modifier of biomass, temperature, pH, initial cobalt concentration, and the dosage of biomass. Each experiment was carried out in 250 ml Erlenmeyer flasks containing 100mL Co(NO<sub>3</sub>)<sub>2</sub>.6H<sub>2</sub>O solution with known initial cobalt concentration which was mixed with different amount of biomass. To adjust the temperature,

Gallenkamp shaker was used, enabling us to set the temperature of the solution. The flasks were shaken at 150 rpm for a period contact time of 90min. The pH of initial solution was adjusted to the desired values with diluted or concentrated HNO<sub>3</sub> and NaOH solutions before mixing with the biosorbent. Cobalt solution was filtrated through Whatman filter paper. Filtered samples were analyzed for residual cobalt ion concentration.

### 3-5. Experimental design

In this study, the effects of operating parameters were optimized using response surface methodology (RSM). RSM is essentially a particular set of mathematical and statistical methods for designing experiments, building models, evaluating the effects of variables, estimating interaction between variables and searching optimum conditions of variables to predict targeted responses [15, 16]. The application of statistical experimental design techniques in sorption processes could result in improved product yields, reduced process variability, as well as reduced development time and overall costs [20].

In this study, central composite design (CCD) was used for the RSM in the experimental design which is well suited for fitting a quadratic surface and usually works well for the process optimization [15, 16]. The CCD consisted of 2<sup>k</sup> factorial runs with 2k axial runs and r center runs. The center points were used to evaluate the experimental error and the reproducibility of the data. The independent variables were coded to the (-1,1) interval where the low and high levels were coded as -1 and +1, respectively. The axial points were located at (±α, 0, 0), (0, ±α, 0) and (0, 0, ±α) where α is

the distance of the axial point from center and makes the design rotatable[5]. In the study, α was fixed at 2.0.

Therefore, central composite design with four numeric factors (initial cobalt (II) concentration: 50–300 mg/l, biosorbent concentration: 0.1–0.5 g/100ml, initial pH: 2.5–7.0 and temperature: 15–45 °C) and one categorical factor at 2 levels (natural *Sargassum sp* and Mg- pretreated *Sargassum sp.* algae) was applied using Design-Expert version 7 (Stat-Ease Inc., Minneapolis, USA) as shown in Table 1. Performance of the process was evaluated by analyzing the response of biosorbent for cobalt (II) ions. The responses were biosorption capacity of natural and modified algae, y<sub>1</sub> and y<sub>2</sub> respectively.

In the optimization process, the responses can be simply related to chosen factors by linear or quadratic models. A quadratic model, which also includes the linear model, is given as:

$$y = b_0 + \sum_{i=1}^k b_i x_i + \sum_{i=1}^k b_{ii} x_i^2 + \sum_{i=1}^k \sum_{j=1}^k b_{ij} x_i x_j \quad (10)$$

where  $b_0$ ,  $b_i$ ,  $b_{ii}$ ,  $b_{ij}$  are a constant, a linear coefficient, a square coefficient and an interaction coefficient, respectively [15, 21, 22].

In this work a central composite design was used to estimate the effect of five independent factors (temperature,  $x_1$ ; initial concentration of cobalt (II),  $x_2$ ; biosorbent dosage,  $x_3$ ; initial pH,  $x_4$ ; natural brown algae and Mg- pretreated brown algae,  $x_5$ ) on the biosorption capacities ( $y_1$ ,  $y_2$ ) as the responses were processed for Eq. (10) including analysis of variance (ANOVA) to obtain the interaction between the process

variables and the responses. The quality of the fit of polynomial model was expressed by the coefficient of determination  $R^2$  and  $R_{adj}^2$ . The statistical significance was checked with adequate precision ratio and F-test [5]. The second-order polynomial models were represented as response, whilst keeping third variable constant.

#### 4. Results and discussion

##### 4-1. Statistical analysis

In order to determine the significant terms 56 runs were statistically analyzed with CCD design. The p-values were used as a tool to check the significance of every coefficient. The smaller the magnitude of p, the more significant the corresponding coefficient is. Values of p less than 0.05 indicate model terms are significant. In this case all five factors, AB (T,  $C_0$ ), BC ( $C_0$ ,  $M_{ads}$ ), BD ( $C_0$ ,

pH) and BE ( $C_0$ , Type of biosorbent) interaction terms are highly significant terms ( $p < 0.05$ ). By eliminating the insignificant model terms automatically, the resulting ANOVA for the reduced quadratic model is summarized in Table 2. The final responses for the biosorption capacity of untreated and treated biomasses for cobalt (II) ions were obtained in Eqs. (11) and (12), respectively.

$$y_1 = 27.01193 - 0.039876 x_1 + 0.21785 x_2 - 109.62840 x_3 + 2.06483 x_4 + 9.15733 E - 004 x_1 x_2 - 0.13322 x_2 x_3 + 0.010372 x_2 x_4 - 3.85354 E - 004 x_2^2 + 98.69344 x_3^2 - 0.35106 x_4^2 \quad (11)$$

$$y_2 = 26.75589 - 0.039876 x_1 + 0.24190 x_2 - 109.62840 x_3 + 2.06483 x_4 + 9.15733 E - 004 x_1 x_2 - 0.13322 x_2 x_3 + 0.010372 x_2 x_4 - 3.85354 E - 004 x_2^2 + 98.69344 x_3^2 - 0.35106 x_4^2 \quad (12)$$

**Table 1.** The experimental range and levels of the control factors in the CCD.

Variable	Low axial (- $\alpha$ )	Low factorial (-1)	Center	High factorial (+1)	High axial (+ $\alpha$ )
$x_1$ (A): Temperature ( $^{\circ}$ C)	15	23	30	38	45
$x_2$ (B): Initial cobalt (II) concentration (mg/l)	50	113	175	238	300
$x_3$ (C): Biomass dose (g/100ml)	0.1	0.2	0.3	0.4	0.5
$x_4$ (D): pH	2.5	3.6	4.8	5.9	7
	Level 1		Level 2		
$x_5$ (E): Type of biosorbent	Natural <i>Sargassum sp.</i>		Mg- pretreated <i>Sargassum sp.</i>		

\*  $\alpha=2$

**Table 2.** ANOVA for the response surface reduced quadratic model for cobalt biosorption .

Source	Sum of squares	df	Mean square	F value	p-value
Model	6424.25	12	535.35	386.89	< 0.0001
A-T	39.13	1	39.13	28.28	< 0.0001
B-C <sub>0</sub>	3255.71	1	3255.71	2352.86	< 0.0001
C-Mads	2609.04	1	2609.04	1885.52	< 0.0001
D-pH	18.03	1	18.03	13.03	0.0008
E-E	218.63	1	218.63	158.00	< 0.0001
AB	5.90	1	5.90	4.26	0.0451
BC	22.18	1	22.18	16.03	0.0002
BD	17.02	1	17.02	12.30	0.0011
BE	27.10	1	27.10	19.59	< 0.0001
B <sup>2</sup>	116.01	1	116.01	83.84	< 0.0001
C <sup>2</sup>	49.87	1	49.87	36.04	< 0.0001
D <sup>2</sup>	10.11	1	10.11	7.30	0.0098
Residual	59.50	43	1.38		
Lack of Fit	52.89	37	1.43	1.30	0.4017
Pure Error	6.61	6	1.10		
Cor Total	6483.75	55			

Non-significant value lack of fit shows the validity of the quadratic model for biosorption by *Sargassum sp.* biomass. The obtained values for the correlation between the actual and predicted response ( $R^2$ ) and coefficient adjusted  $R^2$  are 0.9908 and 0.9883, respectively. These values, which are closer to 1.0 are in reasonable agreement for cobalt (II), indicating the better fitness of the reduced quadratic model in the experimental data. Adequate precision measures the signal to noise ratio and a ratio greater than 4 is


desirable. The ratio of 77.361 indicates an adequate signal. This model can be used to navigate the design space. The CV value was found to be 3.13 for biosorption capacity. Since CV is a measure expressing standard deviation as a percentage of the mean, the smaller values of CV give better reproducibility. In general, a CV higher than 10 indicates that variation in the mean value is high and does not satisfactorily develop an adequate response model [16]. Data in Table 3 confirmed these results.

**Table 3.** Statistical results of the ANOVA for reduced quadratic model.

$R^2$	0.9908
Adjusted $R^2$	0.9883
C.V %	3.13
Adequate Precision	77.361




The predicted values calculated from the statistical model versus the actual response values obtained by the experiment were shown in Fig. 1. It approves the suitability of the model since all the points are located around the diagonal line. The normal probability and studentized residuals plot is shown in Fig. 2 for the biosorption capacity of *Sargassum Sp.* for cobalt (II) ions. In Fig. 2, residuals show how well the model satisfies the assumptions of the analysis of variance (ANOVA) where the studentized residuals measure the number of standard deviations separating the actual and predicted values. Fig. 2 shows that no response transformation needed nor was there an apparent problem with normality.



**Figure 1.** Scatter graph of the predicted response values versus the actual response values for the biosorption capacity of biomass for cobalt.

#### 4-2. Statistical optimization

The aim of the process was to maximize biosorption capacity of biosorbent for cobalt (II) ions and find optimized conditions. The optimum values for temperature, initial cobalt (II) concentration, biosorbent dosage and pH were estimated to be 45°C, 300 mg/l, 0.1 g/100ml, and 7.0, respectively. According to the optimum conditions, the best biosorbent which had the maximum metal uptake (80.5 mg/g) was Mg(NO<sub>3</sub>)<sub>2</sub>.6H<sub>2</sub>O-treated biomass (Mg-treated biomass). The results for optimized parameters by Design-Expert software are presented in Table 4.



**Figure 2.** Normal plot of studentized residuals versus normal % probability for bearing the experiment for cobalt ions.

**Table 4.** Optimum values of the process parameters for maximum biosorption capacity.

Parameters	T(°C)	C <sub>0</sub> (mg/l)	M <sub>ads</sub> (g)	pH	Type of biosorbent
Value	45	300	0.1	7	Mg-treated biomass

The software calculated the expected responses and associated confidence intervals based on the prediction equations (Eqs. (11) and (12)). The 95% confidence interval (C.I.) is the range in which the process average was expected to fall 95% of the time, while the 95% prediction interval (P.I.) is the range in which it was expected that any individual value would fall 95% of the time[23]. As shown in Table 5, the P.I. is larger (a wider spread) than the C.I. since more scatter in individual values is expected than in averages. In this Table, SE mean refers to the standard deviation associated with the prediction of an average value at the selected component-levels, while the standard deviation associated with SEpred. indicates the prediction of an individual observation at the selected factor levels.


#### 4-3. Effect of pretreatment on biosorption capacity and sorption mechanism

The chemically modified biosorbents increase the stability of the biosorbent material and enhance the biosorbent properties[24]. The interactive effect of initial cobalt (II) concentration and type of biomass on the cobalt uptake of biomass holding other variables at their central values is shown in Fig. 3. The capacities of treated *Sargassum sp.* and untreated *Sargassum sp.* for cobalt ion at different concentration noted that the treatment of  $Mg(NO_3)_2$  had enhanced the adsorption capacity. At the sorption of cobalt, the adsorption capacity for untreated *Sargassum sp.* and treated *sargassum sp.*


increased from 15.43 to 45.49 mg/g and 16.37 to 52.45 mg/g with the increase of the initial concentration in 50–300 mg/l, respectively. According to the Table 2, the interaction between initial cobalt (II) concentration and type of biomass (BE) was highly significant ( $p < 0.0001$ ). EDX spectra taken in spot profile mode showed the enhancement of magnesium peaks on EDX spectra of biomass after treatment compared with biosorbent before Mg-treatment (Fig. 4a, b) and also confirmed the presence of cobalt (Fig.4c), suggesting bonding of metal cations onto algae surface by interactions with negatively charged functional groups. Additionally, for the samples after biosorption, we observed that the peaks of Mg on EDX spectra of biosorbent after metal sorption were omitted (Fig. 4c).The amount of magnesium and cobalt of the untreated *Sargassum sp.*, Mg-treated biomass and cobalt-loaded biomass was determined using anEDX spectrometer, given in Table 6. Our data suggest that  $Co^{2+}$  ions have replaced  $Mg^{2+}$  ions from biosorbent surface and ion exchange mechanism participates in cobalt biosorption. The same conclusions were also postulated by others [25-28]. But, due to the complexity of biomaterials, maybe other mechanisms such as coordination and chelation of metals, adsorption and surface complexation of cations with exposed functional groups on the biomass act simultaneously, to varying degrees, depending on the biosorbent and the solution chemistry[8, 29-31].

**Table 5.** Point prediction of the responses at the optimal conditions.

Response	Prediction	SE Mean	95%C.I. low	95% C.I. high	SE pred.	95% P.I.low	95% P.I. high
q (mg/g)	80.2094	1.94	76.29	84.13	2.27	75.63	84.79



**Figure 3.** Interaction plot showing effect of initial cobalt (II) concentration and type of biomass on the cobalt uptake of biomass holding other variables at their central values.



**Figure 4.** EDX analysis of *Sargassum sp.* before (a) and after (b) Mg treatment, and after (c) cobalt (II) sorption. Au peak comes from sample coating.


**Table 6.** The amount of magnesium and cobalt of the biomass.

Type of biomass	Element [wt. %]	
	Magnesium	Cobalt
Untreated <i>Sargassum sp.</i>	1.91	0.0
Mg-treated <i>Sargassum sp.</i>	4.19	0.0
cobalt-loaded <i>Sargassum sp.</i>	0.0	11.04

#### 4-4. Effect of operating parameters on biosorption



As shown in Table 2, the results indicated that all linear and quadratic parameters and mutual interaction between temperature and initial cobalt (II) concentration (AB), initial cobalt (II) concentration and biomass dosage (BC) and initial cobalt (II) concentration and pH (BD) were significant ( $p < 0.0001$ ,  $p < 0.05$ ). The response surface plots (three-dimensional graphs and contour plots) presented in Figs. 5-7 were generated by varying two factors while keeping the others constant in center values.

Adsorption solution pH influences both cell surface metal binding sites and metal chemistry in water [14]. Cobalt predominantly exists as  $\text{Co}^{2+}$  cation within a broad pH range from 2.5 to 7.0. The concentration of  $\text{Co}^{2+}$  starts to decrease at pH 7.5. The precipitation of cobalt (formation of  $\text{Co}(\text{OH})_2$  form) started at pH 7.5. Therefore, the pH range studied was between 2.5 and 7.0, pH values above 7 were not studied because cobalt removal by hydroxide formation begins to interfere with cobalt removal by biosorption and it will be difficult to distinguish between cobalt biosorption and precipitation. Fig. 5 shows the effects of initial cobalt (II) concentration and pH (BD)





**Figure 5.** Three-dimensional plots showing effects of initial cobalt (II) concentration and pH (BD) on the biosorption capacity for Mg-treated biomass holding other variables at their central values.

on the biosorption capacity for Mg-treated biomass. Cobalt uptake significantly increases with metal concentration but a smaller increase with pH is observed. According to Fig. 5, the biosorption capacity at pH 7.0 and pH 2.5 was 54.73 and 46.44 mg/g, respectively, and the  $\text{Co}^{2+}$  concentration was 300 mg/l while the other parameters are in the center values ( $T = 30^\circ\text{C}$  and  $M_{\text{ads}} = 0.3$ ). The data show the biosorption uptake did not vary significantly with initial pH, indicating that the biosorbent



**Figure 6.** Three-dimensional plots showing effects of temperature and initial cobalt (II) concentration (AB) on the cobalt uptake of Mg-treated biomass holding other variables at their central values.



**Figure 7.** Three-dimensional plots showing effects of initial cobalt (II) concentration and biomass dosage (BC) holding the temperature at 30°C and pH 4.8.

can be used with comparable effectiveness in the pH range 2.5 to 7.0. The biosorption-capacity decreases with decreasing pH, which is explained by the greater affinity of the proton by the binding sites at the algal particles surface when compared with the metal ion. Conversely, at higher pH where cobalt concentration is much greater than  $H^+$  concentration, the metal uptake increases. Similar results have been obtained by other authors [5, 29] who found an optimal pH range of 4–7 [5, 25, 29, 30]. At this pH range some functional groups in the algae surface, e.g. carboxyl and sulfate, become negatively charged, increasing the electrostatic interactions with the metal cation [5].

The biosorption is highly dependent on the amount of metal initially present in solution, increasing with the initial concentration. For example, the biosorption capacity changes about +41.79 mg/g by increasing initial  $Co^{2+}$  concentration from 50 to 300 mg/l at pH 7. Higher concentration of cobalt ions in the solution causes the active sites of *Sargassum sp.* to be surrounded by more cobalt ions, and biosorption would be carried out more sufficiently. Similar results are also reported

by researchers for a variety of adsorbate-adsorbent systems [3, 32]. An increase of metal uptake by increasing initial metal ion concentration is a result of the increase in the driving force of the concentration gradient, rather than increase in the initial metal ion concentration [33]. The initial concentration provides an important driving force to overcome all mass transfer resistance of cobalt between the aqueous and solid phases. The effect of temperature on the cobalt uptake of Mg-treated biomass holding other variables at their central values is shown in Fig. 6. The biosorption capacity of treated *Sargassum sp.* increased with the increase in biosorption temperature and initial cobalt (II) concentration. The maximum biosorption capacity occurred at 45°C. However, temperature variation between 15 and 45°C slightly increases the metal uptake capacity from 38.63 to 42.24 mg/g, respectively. The uptake capacity was calculated at pH 4.8, 175 mg/l initial cobalt (II) concentration and 0.3 g biomass dosage. These results indicate the negligible interaction effect of these two parameters on biosorption capacity which confirmed the result of ANOVA as shown in Table 2 (p value = 0.0451). The increment of sorption capacity with increasing temperature suggests an endothermic nature of the sorption of cobalt by this dried biomass. Several studies have indicated the same effect of temperature [13, 34-36]. Higher temperature causes rupture that enhances the number of available active sites, surface activity, kinetic energy of the solute, and affinity of sites for metal ions which lead to an increase in the probability of the collision between active surface binding sites and metal ions and a decrease in the thickness of the boundary layer surrounding the biosorbent [13, 35-37]. However, the temperature higher than 55°C caused a change in the texture of the biomass and thus

reduced its sorption capacity [14, 38]. In addition, the decrease of biosorption efficiency with the increase of temperature is due, most probably, to the desorption tendency of heavy metals from biosorbent surface. [39] Therefore, it depends on the type of metal and adsorbent. That is the reason for having different behavior of heavy metal uptake with temperature [38].

The effects of biomass dosage (BC) can be inferred from the response plot (Fig. 7), holding the temperature at 30°C and pH at 4.8. The biosorption capacity was decreased from 59.13 to 29.64 mg/g for the  $\text{Co}^{2+}$  ions with increasing the biosorbent dosage from 0.1 to 0.5 g, respectively. The uptake capacity was calculated at pH 4.8 and initial metal ion concentration 175 mg/l. At higher biosorbent dose the uptake capacity is low, perhaps due to the unsaturation of biosorption sites through the biosorption reaction and due to the particle interaction, such as aggregation, resulted from high biosorbent concentration [40]. Such aggregation would lead to decrease in total surface area of the biosorbent and an increase in diffusional path length [41].

#### 4-5. Confirmatory experiments

For validation of the statistical model, two experiments were performed under optimal conditions. Table 7 presents the experimental results under the optimal condition compared with the simulated values from the proposed model (Eq. (12)). The experimental responses for biosorption capacity were 81.11 and 79.98 mg/g which indicate that the experimental values are very close to the predicted values and hence, the model is successful in predicting the responses.

**Table 7.** Optimum condition verification and additional experiments.

Run	T(°C)	C <sub>0</sub> (mg/l)	M <sub>ads</sub> (g)	pH	Type of biosorbent	Response (maximized Co <sup>2+</sup> uptake)	
						q <sub>pre</sub> (mg/g)	q <sub>exp</sub> (mg/g)
1	45	300	0.1	7	Mg-treated biomass	80.21	79.98
2	45	300	0.1	7	Mg-treated biomass	80.21	81.11


#### 4-6. Determination of dynamic constants at optimized condition

In order to obtain the biosorption kinetics constants of cobalt ions on Mg-treated biomass, two kinetics models were studied at optimized conditions determined by Design-Expert software (T=45°C, C<sub>0</sub>=300 ppm, pH 7 and M<sub>ads</sub>= 0.1 g). According to the results, the theoretical q<sub>e(cal)</sub> values calculated from the pseudo-first-order model did not give reasonable values with regard to the experimental uptake ones, q<sub>e(exp)</sub>. Further, the correlation coefficient (R<sup>2</sup>=0.7224) is less than 0.99, suggesting that the present adsorption system does not follow pseudo-first-order process. Similar results are also reported by other researchers[1, 17, 42, 43].

The plots between t/q<sub>t</sub> and t for the pseudo-second-order model are shown in Fig. 8. As can be seen from Table 8, the theoretical q<sub>e(cal)</sub> values agree well with the experimental uptake values, q<sub>e(exp)</sub>. Further, the correlation coefficient (R<sup>2</sup>) for the pseudo-second-order kinetic model was 0.999, suggesting that the present adsorption system can be described more favorably by pseudo-second-order process.

#### 4-7. Isotherm at optimized condition

The biosorption isotherm for the sorption on cobalt on Mg-treated biomass was obtained at different temperature and constant



**Figure 8.** The linearized pseudo-order biosorption dynamics of cobalt.

**Table 8.** Biosorption kinetics constants of cobalt ions on Mg-treated biomass for pseudo-second-order model at optimized conditions (T=45 °C, C<sub>0</sub>=300 ppm, pH 7 and M<sub>ads</sub>= 0.1 g).

q <sub>e (exp)</sub> (mg/g)	k <sub>2</sub> (g/mg min)	q <sub>e (cal)</sub>	R <sup>2</sup>
82.789	0.058564	82.645	0.999

optimized conditions determined by Design-Expert software (pH 7 and M<sub>ads</sub>= 0.1). The results of the adsorption isotherms for the cobalt ions at different temperatures are presented in Table 9. The good correlation coefficients values (R<sup>2</sup>) showed that Freundlich model is more suitable than Langmuir for adsorption equilibrium of cobalt. The better fit of equilibrium data to the Freundlich equation refers the multilayer sorption of the cobalt ions on the biosorbent [41]. However, some studies have been

**Table 9.** Adsorption isotherm parameters for Co (II) biosorption on Mg-treated *Sargassum sp.*

T (°C)	Langmuir constants			Freundlich constants		
	q <sub>m</sub> (mg/g)	b (l/mg)	R <sup>2</sup>	k	n	R <sup>2</sup>
15	76.33588	0.024472	0.9709	7.485141	2.398657	0.9987
23	78.74016	0.023523	0.9719	7.247697	2.336449	0.9988
30	80.64516	0.02274	0.9728	7.048553	2.285192	0.9988
38	83.33333	0.021779	0.9738	6.83282	2.230152	0.9989
45	85.47009	0.021051	0.9748	6.652732	2.184837	0.999


described better with the Langmuir isotherm [27, 29, 39]. Some authors explain that Langmuir isotherm corresponds to a dominant ion exchange mechanism while the Freundlich isotherm shows adsorption-complexation reactions taking place in the adsorption process [44]. The result in Table 9 shows that n values related to Freundlich isotherm are greater than unity representing favorable biosorption conditions and formation of relatively stronger bonds between adsorbent and adsorbate [3, 5].

#### 4-8. Determination of thermodynamic constants of biosorption

The calculated thermodynamic parameter Gibbs free energy change,  $\Delta G^\circ$ , for the biosorption of cobalt by *Sargassum sp.* algae was calculated using Eq. 7 and 9 (Table 10). According to Eq. 9, the enthalpy change,  $\Delta H^\circ$ , and the entropy change,  $\Delta S^\circ$ , were calculated from the plot of  $\ln K$  versus  $1/T$  ( $R^2=0.999$ ) to give the values 535.76 J/mol and 3.24 J/molK, respectively (Fig. 9). The negative value of  $\Delta G^\circ$  indicates the feasibility of the process and indicates the spontaneous nature of the adsorption. The  $\Delta G^\circ$  value is more negative with increasing temperature, which suggests that higher temperature makes the adsorption easier. In

general, it is of note that up to  $-20$  kJ/mol is consistent with electrostatic interaction between charged molecules and surface indicative of physisorption while more negative than  $-40$  kJ/mol involves chemisorption and values between  $-20$  and  $-40$  kJ/mol indicate that both physisorption and chemisorption were responsible for adsorption[17].

For the adsorption of cobalt ions on to the *Sargassum sp.* biomass, the order of magnitude of the  $\Delta G^\circ$  value indicates a physical mechanism. However, the pseudo-second-order kinetics shows chemisorption mechanism for this system. Therefore, the nature of the adsorption process cannot be clearly specified thermodynamically, and it might only be considered as chemisorption in nature enhanced by physical effects.

**Figure 9.** Thermodynamic relationship between the Gibbs free energy and temperature.



In our study, the positive value of  $\Delta H^\circ$  confirms the endothermic adsorption nature of the process. The positive value of  $\Delta S^\circ$  indicates the affinity of the adsorbent for cobalt and also confirms the increased randomness at the solid-solution interface

during biosorption [2, 17]. Value of the  $\Delta G^\circ$ ,  $\Delta H^\circ$  and  $\Delta S^\circ$  from the present study was compared with other metal adsorption reported in previous studies and are compiled in Table 11.

**Table 10.** Thermodynamic parameters and corresponding correlation coefficient for Co (II) biosorption on Mg-treated *Sargassum sp.*

Temperature (°C)	Thermodynamic parameters and corresponding R <sup>2</sup> value			
	$\Delta G^\circ$ (J/mol)	$\Delta H^\circ$ (J/mol)	$\Delta S^\circ$ (J/mol K)	R <sup>2</sup>
15	-397.935862	535.7625	3.239966	0.999
23	-422.706613			
30	-445.18927			
38	-471.420463			
45	-495.217534			

**Table 11.** Comparison of thermodynamic parameters with other metals adsorption.

Heavy metal ion	Sorbent	Temperature (°C)	$\Delta G^\circ$ (kJ mol <sup>-1</sup> )	$\Delta H^\circ$ (kJ mol <sup>-1</sup> )	$\Delta S^\circ$ (J mol <sup>-1</sup> K <sup>-1</sup> )	Reference
Pb <sup>2+</sup>	Peanut shells	20, 30, 40	-24.25 to -25.00	-16.68	25.16	[18]
Pb <sup>2+</sup>	Sargassumilicifolium	20, 25, 30	-2.6 to -3.6	0.027	0.102	[13]
Sr <sup>2+</sup>	Activated carbon	20, 30, 40, 60	-12.61 to -20.44	44.77	195.847	[45]
Cr (VI)	Sargassummuticum	20, 50	-7.3 to -8.7	7.1	49.1	[36]
Pb <sup>2+</sup> Ni <sup>2+</sup> Cd <sup>2+</sup>	Orange peel	30	-3.77, -4.22, -4.99	-	-	[46]
U	Padina sp.	10, 20, 30, 40	-6.549 to -5.914	-12.552	-21.408	[17]
Cu <sup>2+</sup> Cd <sup>2+</sup> Pb <sup>2+</sup>	Caulerpalentillifera	20, 30, 40	-16.4 to -18.5 -13.4 to -17.2 -17.9 to -19.7	13.9 42.3 8.28	103 191 89.3	[35]
Ni <sup>2+</sup>	S. muticum G. caudate	20, 30, 50	-	-29.7 -16.0	-90.3 -47.7	[47]
Co <sup>2+</sup>	Lemon peel	25, 45	-37.47, -38.56	-21.20	54.61	[1]
Sr <sup>2+</sup>	Activated carbon	20, 30,40,60	-36.61 to -41.75	0.036.88	125.44	[48]
Co <sup>2+</sup>	<i>Sargassum Sp.</i>	15, 23, 30, 38, 45	-0.397 to -0.495	0.535	3.24	This work

## 5. Conclusions

In this study, central composite design (CCD) was used for the RSM in the experimental design and proved to be an efficient method for testing the effect of operating conditions and their interactions on cobalt uptake by the brown algae *Sargassum sp.* The interactive effects of five independent factors: initial pH of solution, initial concentration cobalt (II), temperature, biosorbent dosage, and biomass treatment on the biosorption capacities were estimated. A reduced quadratic model was obtained to predict the biosorption capacity. ANOVA results confirmed that there was significant agreement between the model and experimental data. The optimum biosorption conditions were determined as Mg-treated biomass, initial pH 7.0, temperature 45°C, biosorbent dosage 0.1 g/100ml and initial cobalt (II) ion concentration 300 mg/l. At optimum biosorption conditions, the biosorption capacity of Mg-treated biomass for cobalt (II) ions was found to be 80.55 mg/g. The results are best fitted by the Freundlich model. The biosorption of cobalt (II) onto algal biomass followed the pseudo-second-order dynamics well. The calculated thermodynamic parameters ( $\Delta G^\circ$ ,  $\Delta H^\circ$  and  $\Delta S^\circ$ ) showed that the process was feasible, spontaneous and endothermic at the temperature ranges of 15–45°C.

## Acknowledgment

Special thanks to Dr. S.M. Mousavi and M. Jalaiean for their help in accomplishing this work. The authors are grateful to Stat-Ease, Minneapolis, USA, for the provision of the Design-Expert package.

## Nomenclature

$b$	the Langmuir constant (l/mg)
$b_0, b_i$ and $b_{ij}$	linear and quadratic interaction coefficients, $i$ and $j = 1-5$
$C$	metal ion concentration at any time (mg/l)
$C_0$	initial metal ion concentration (mg/l)
$C_{eq}$	residual metal ion concentration at equilibrium (mg/l)
$C.I.$	confidence interval (-)
$C.V$	coefficient variation (-)
$k$	adsorption capacity (-)
$n$	biosorption intensity (-)
$P.I.$	prediction interval (-)
$q$	amount of biosorbed metal per g of biosorbent at any time (mg/g)
$q_{eq}$	the amount of biosorbed metal per unit weight of biosorbent at equilibrium (mg/g)
$q_{exp}$	experimental amount of biosorbed metal per unit weight of biosorbent at equilibrium (mg/g)
$q_m$	maximum capacity of biosorbent
$q_{pre}$	predicted amount of biosorbed metal per unit weight of biosorbent at equilibrium by software (mg/g)
$R^2$	correlation coefficient
$R^2_{adj}$	adjusted correlation coefficient
$M_{ads}$	amount of biosorbent (g)
$t$	time (min)
$x_i$	independent variable, $i=1-5$
$y$	response (-)
$V$	volume of the solution (l)
$T$	Solution temperature (°C)

## References

- [1] Bhatnagar, A., Minocha, A.K. and Sillanpää, M., "Adsorptive removal of cobalt from aqueous solution by utilizing lemon peel as biosorbent", *Biochem. Eng. J.*, 48(2), 181, (2010).
- [2] Pahlavanzadeh, H., Keshtkar, A.R., Safdari, J. and Abadi, Z., "Biosorption of nickel(II) from aqueous solution by brown algae: Equilibrium, dynamic and thermodynamic studies", *J. Hazard. Mater.*, 175 (1-3), 304, (2010).
- [3] Dahiya, S., Tripathi, R.M. and Hegde, A.G., "Biosorption of heavy metals and radionuclide from aqueous solutions by pre-treated arca shell biomass", *J. Hazard. Mater.*, 150(2), 376, (2008).
- [4] Kalavathy, M.H., Karthikeyan, T., Rajgopal, S. and Miranda, L.R., "Kinetic and isotherm studies of Cu(II) adsorption onto H<sub>3</sub>PO<sub>4</sub>-activated rubber wood sawdust", *J. Colloid. Interf. Sci.*, 292(2), 354, (2005).
- [5] Özer, A., Gürbüz, G., Çalimli, A. and Körbahti, B.K., "Investigation of nickel(II) biosorption on *Enteromorpha prolifera*: Optimization using response surface analysis", *J. Hazard. Mater.*, 152(2), 778, (2008).
- [6] Gupta, V.K. and Rastogi, A. "Biosorption of lead(II) from aqueous solutions by non-living algal biomass *Oedogonium sp.* and *Nostoc sp.*—A comparative study", *Colloids Surf. B.*, 64(2), 170, (2008).
- [7] Romera, E., González, F., Ballester, A., Blázquez, M. L. and Muñoz, J.A., "Comparative study of biosorption of heavy metals using different types of algae", *Bioresour. Technol.*, 98(17), 3344, (2007).
- [8] Davis, T.A., Volesky, B. and Mucci, A., "A review of the biochemistry of heavy metal biosorption by brown algae", *Water Res.*, 37(18), 431, (2003).
- [9] Kleinubing S.J., da Silva, E.A., da Silva, M.G.C. and Guibal, E., "Equilibrium of Cu(II) and Ni(II) biosorption by marine alga *Sargassum filipendula* in a dynamic system: Competitiveness and selectivity", *Bioresour. Technol.*, 102, 4610, (2011).
- [10] Haug, A., "The affinity of some divalent metals to different types of alginates", *Acta Chem. Scand.*, 15, 1794, (1961).
- [11] Papageorgiou, S.K., Katsaros, F.K., Kouvelos, E.P., Nolan, J.W., Le Deit, H. and Kanellopoulos, N.K., "Heavy metal sorption by calcium alginate beads from *Laminaria digitata*", *J. Hazard. Mater.*, 137, 1765, (2006).
- [12] Demim, S., Drouiche, N., Aouabed, A., Benayad, T., Couderchet, M. and Semsari, S. "Study of heavy metal removal from heavy metal mixture using the CCD method", *J. Ind. Eng. Chem.*, 20(2), 512, (2014).
- [13] Tabaraki, R., Nateghi, A. and Ahmady-Asbchin, S., "Biosorption of lead (II) ions on *Sargassum ilicifolium*: Application of response surface methodology", *Intl. biodeter. Biodegr.*, 93, 145, (2014).
- [14] Amini, M., Younesi, H., Bahramifar, N., Lorestani, A.A.Z., Ghorbani, F., Daneshi, A. and Sharifzadeh, M., "Application of response surface

- methodology for optimization of lead biosorption in an aqueous solution by *Aspergillus niger*", *J. Hazard. Mater.*, 154(1–3), 694, (2008).
- [15] Montgomery, D.C., *Design and analysis of experiments*, John Wiley & Sons Co., New York, USA, (2001).
- [16] Myers, R.H. and Montgomery, D.C., *Response surface methodology: Process and product optimization using designed experiments*, John Wiley & Sons Co., New York, USA, (2002).
- [17] Khani, M., "Uranium biosorption by *Padina sp.* algae biomass: Kinetics and thermodynamics", *Environ. Sci. Pollut. Res.*, 18(9), 1593, (2011).
- [18] Taşar, Ş., Kaya, F. and Özer, A., "Biosorption of lead(II) ions from aqueous solution by peanut shells: Equilibrium, thermodynamic and kinetic studies", *J. Environ. Chem. Eng.*, 2(2), 1018, (2014).
- [19] Tunç, Ö., Tanacı, H. and Aksu, Z., "Potential use of cotton plant wastes for the removal of Remazol Black B reactive dye", *J. Hazard. Mater.*, 163(1), 187, (2009).
- [20] Annadurai, G., Juang, R.S. and Lee, D.J., "Adsorption of heavy metals from water using banana and orange peels", *Water Sci. Technol.*, 47, 185, (2002).
- [21] Goel, J., Kadirvelu, K., Rajagopal, C. and Garg, V.K., "Removal of Lead(II) from aqueous solution by adsorption on carbon aerogel using a response surface methodological approach", *Ind. Eng. Chem. Res.*, 44(7), 1987, (2005).
- [22] Amani, T., Nosrati, M., Mousavi, S.M. and Kermanshahi, R.K., "Study of syntrophic anaerobic digestion of volatile fatty acids using enriched cultures at mesophilic conditions", *Intern. J. Environ. Sci. Technol.*, 8, 83, (2011).
- [23] Mafi Gholami, R., Mousavi, S.M. and Borghei, S.M., "Process optimization and modeling of heavy metals extraction from a molybdenum rich spent catalyst by *Aspergillus niger* using response surface methodology", *J. Ind. Eng. Chem.*, 18(1), 218, (2012).
- [24] Montazer-Rahmati, M.M., Rabbani, P., Abdolali, A. and Keshtkar, A.R., "Kinetics and equilibrium studies on biosorption of cadmium, lead, and nickel ions from aqueous solutions by intact and chemically modified brown algae", *J. Hazard. Mater.*, 185(1), 401, (2011).
- [25] Vijayaraghavan, K., Jegan, J., Palanivelu, K. and Velan, M., "Biosorption of cobalt(II) and nickel(II) by seaweeds: Batch and column studies", *Sep. Purif. Technol.*, 44(1), 53, (2005).
- [26] Bishnoi, N.R., Kumar, R., Kumar, S. and Rani, S., "Biosorption of Cr(III) from aqueous solution using algal biomass *spirogyra spp.*", *J. Hazard. Mater.*, 145(1–2), 142, (2007).
- [27] Hatzikioseyan, A., Tsezos, M. and Mavituna, F., "Application of simplified rapid equilibrium models in simulating experimental breakthrough curves from fixed bed biosorption reactors", *Hydrometallurgy*, 59(2–3), 395, (2001).
- [28] Kuyucak, N. and Volesky, B., "Accumulation of cobalt by marine alga", *Biotechnol. Bioeng.*, 33(7), 809, (1989).

- [29] Suhasini, I.P., Sriram, G., Asolekar, S.R. and Sureshkumar, G.K., "Biosorptive removal and recovery of cobalt from aqueous systems", *Process. Biochem.*, 34(3), 239, (1999).
- [30] Marešová, J., Pipiška, M., Rozložník, M., Horník, M., Remenárová, L. and Augustín, J., "Cobalt and strontium sorption by moss biosorbent: Modeling of single and binary metal systems", *Desalination*, 266(1–3), 134, (2011).
- [31] Sheng, P.X., Ting, Y.P., Chen, J.P. and Hong, L., "Sorption of lead, copper, cadmium, zinc, and nickel by marine algal biomass: Characterization of biosorptive capacity and investigation of mechanisms", *J. Colloid. Interf. Sci.*, 275(1), 131, (2004).
- [32] Ahmadpour, A., Tahmasbi, M., Bastami, T.R. and Besharati, J.A., "Rapid removal of cobalt ion from aqueous solutions by almond green hull", *J. Hazard. Mater.*, 166(2–3), 925, (2009).
- [33] Ghorbani, F., Younesi, H., Ghasempouri, S.M., Zinatizadeh, A.A., Amini, M. and Daneshi, A., "Application of response surface methodology for optimization of cadmium biosorption in an aqueous solution by *Saccharomyces cerevisiae*", *Chem. Eng. J.*, 145(2), 267, (2008).
- [34] Özcan, A.S., Tunali, S., Akar, T. and Özcan, A., "Biosorption of lead(II) ions onto waste biomass of *Phaseolus vulgaris* L.: Estimation of the equilibrium, kinetic and thermodynamic parameters", *Desalination*, 244, 188, (2009).
- [35] Apiratikul, R., Madacha, M. and Pavasant, P., "Kinetic and mass transfer analyses of metal biosorption by *Caulerpalentil-lifera*", *Desalination*, 278, 303, (2011).
- [36] Bermúdez, Y.A., Rico, I.L., Guibal, R.E., de Hoces, M.C. and Martín-Lara, M.A., "Biosorption of hexavalent chromium from aqueous solution by *Sargassummuticum* brown alga: Application of statistical design for process optimization", *Chem. Eng. J.*, 183, 68, (2012).
- [37] Freitas, O., Delerue-Matos, C. and Boaventura, R., "Optimization of Cu(II) biosorption onto *Ascophyllumnodosum* by factorial design methodology", *J. Hazard. Mater.*, 167(1–3), 449, (2009).
- [38] Qaiser, S., Saleemi, A. and Mahmood Ahmad, M., "Heavy metal uptake by agro based waste materials", *Elec. J. Biotechnol.*, 10 (3), 409, (2007).
- [39] Bulgariu, D. and Bulgariu, L., "Equilibrium and kinetics studies of heavy metal ions biosorption on green algae waste biomass", *Bioresour. Technol.*, 103, 489, (2012).
- [40] Özer, A., Gürbüz, G., Çalimli, A. and Körbahti, B.K., "Biosorption of copper(II) ions on *Enteromorpha prolifera*: Application of response surface methodology (RSM)", *Chem. Eng. J.*, 146(3), 377, (2009).
- [41] Singh, R., Chadetrik, R., Kumar, R., Bishnoi, K., Bhatia, D., Kumar, A., Bishnoi, N.R. and Singh, N., "Biosorption optimization of lead(II), cadmium(II) and copper(II) using response surface methodology and applicability in isotherms and thermodynamics modeling", *J. Hazard.*

- Mater.,174(1-3), 623, (2010).
- [42] Cruz, C.C.V., da Costa, A.C.A., Henriques, C.A. and Luna, A.S., "Kinetic modeling and equilibrium studies during cadmium biosorption by dead *Sargassum sp.* biomass", Bioresour. Technol.,91(3), 249, (2004).
- [43] Sun, X.F., Wang, S.G., Liu, X.W., Gong, W.X., Bao, N. and Gao, B.Y., "Competitive biosorption of zinc(II) and cobalt(II) in single- and binary-metal systems by aerobic granules", J. Colloid. Interf. Sci., 324(1-2), 1, (2008).
- [44] Khani, M., "Statistical analysis and isotherm study of uranium biosorption by *Padina sp.* algae biomass", Environ. Sci. Pollut. Res., 18(5), 790, (2011).
- [45] Kacan, E. and Kütahyalı, C., "Adsorption of strontium from aqueous solution using activated carbon produced from textile sewage sludges", J. Anal. Appl. Pyrolysis, 97, 149, (2012)
- [46] Feng, N., Guo, X., Liang, S., Zhu, Y., Liu, J., "Biosorption of heavy metals from aqueous solutions by chemically modified orange peel", J. Hazard. Mater.,185, 49, (2011)
- [47] Bermúdez, Y.G., Rico, I.L. and Bermúdez, O.G., "Nickel biosorption using *Gracilariacaudata* and *Sargassummuticum*", Chem. Eng. J., 166, 122, (2011).
- [48] Chegrouche, S., Mellah, A. and Barkat, M., "Removal of strontium from aqueous solutions by adsorption onto activated carbon: Kinetic and thermodynamic studies", Desalination, 235, 306, (2009).